

G.M. Wysin,\* F.G. Mertens†

\*Kansas State University, Manhattan, KS 66506 USA

†Physics Institute, University of Bayreuth,  
D-8580 Bayreuth, Germany

The dynamics of individual and pairs of vortices in a classical easy-plane Heisenberg spin model is studied. There are two types of vortices possible: in-plane, with small out-of-plane spin components present only at nonzero velocity, and out-of-plane, with large out-of-plane spin components even when at rest. As a result, the two types are governed by different equations of motion when in the presence of neighboring vortices. We review the static spin configurations and the changes due to nonzero velocity. An equation of motion introduced by Thiele and used by Huber will be re-examined. However, that equation may be inadequate to describe vortices in the XY model, due to their zero gyrovectors. An alternative dynamical equation is developed, and effective mass and dissipation tensors are defined. These are relevant for models with spatially anisotropic coupling in combination with easy-plane spin exchange.

## INTRODUCTION

A model for the dynamic correlations of vortices in easy-plane two-dimensional magnets has been presented, that uses the idea of an ideal gas of weakly interacting vortices.<sup>1</sup> Assuming a Boltzmann velocity distribution, and if the velocity-dependent spin field of the vortices is known, then the dynamic structure function  $S^{\alpha\alpha}(\vec{q}, \omega)$  can be determined. At the microscopic level we would like to investigate the time-dependent motion of a single

vortex, to understand how the neighboring vortices cause forces and accelerations, and to have a clear picture of how equilibrium is achieved.

Huber<sup>2,3</sup> has done such an analysis for diffusive motion of so-called "out-of-plane" vortices, ones that possess large out-of-easy-plane spin components. However, it is now realized that there are two type of vortices possible,<sup>4,5</sup> depending on the strength of the easy-plane anisotropy.<sup>6,7</sup> The stable vortices of the XY model, for example, are so-called "planar" vortices that only have small out-of-plane spin components. In that case the equation of motion that was used<sup>3,8</sup> is found to be inapplicable because these planar vortices have a zero gyrovector, to be discussed below. Here we propose an alternative dynamic equation of motion that applies to both types of vortices.

We begin by summarizing the properties of the two types of vortices allowed in the easy-plane anisotropic ferromagnetic Heisenberg model. The derivation of the equation of motion introduced by Thiele,<sup>8</sup> in terms of conserved force densities, will be sketched out, and the breakdown for planar vortices will be discussed. An alternative formalism using a canonical momentum for the vortex is developed. The new equation of motion includes the effects of vortex shape changes that are the result of acceleration. This leads to definition of an effective mass tensor, and, the gyrovector also re-appears. The new equation allows for a consistent description of both types of vortices.

### Anisotropic Heisenberg Ferromagnet

The model system is the nearest neighbor 2D Heisenberg ferromagnet with easy-plane anisotropic exchange, characterized by a parameter  $0 \leq \lambda < 1$ ; the Hamiltonian is

$$\mathbf{H} = -J \sum_{ij} (S_i^x S_j^x + S_i^y S_j^y + \lambda S_i^z S_j^z) . \quad (1)$$

$J$  is an energy scale and the  $\vec{S}_i$  are classical spins with fixed length. The XY model is given when  $\lambda=0$ , the Heisenberg model when  $\lambda=1$ . The spin dynamics is described by the Landau-Lifshitz equation,<sup>9,10</sup>

$$\dot{\vec{S}}_i = \{\vec{S}_i, H\} - \alpha \vec{S}_i \times \dot{\vec{S}}_i = \vec{S}_i \times (\vec{H}_i - \alpha \dot{\vec{S}}_i) \quad (2a)$$

$$\vec{H}_i = J \sum_{(ij)} (S_j^x \hat{x} + S_j^y \hat{y} + \lambda S_j^z \hat{z}) \quad (2b)$$

The sum is only over the neighbors of site  $i$ . A Landau-Gilbert damping term of strength  $\alpha$  has been included. At any given time each spin is instantaneously precessing about the effective

field  $(\vec{H}_i - \alpha \dot{\vec{S}}_i)$ . Initially the vortices will be described in the absence of damping,  $\alpha=0$ , which can be later re-introduced at the phenomenological level.

### Static Vortices

The spins are parametrized in terms of an in-plane angle  $\phi(\vec{x}, t)$  and an out-of-plane angle  $\theta(\vec{x}, t)$  (or we use  $S^z = S \sin\theta$ ),

$$\vec{S}(\vec{x}, t) = S(\cos\theta \cos\phi, \cos\theta \sin\phi, \sin\theta) . \quad (3)$$

Then in a continuum limit including terms up to 2nd order in gradients the equations of motion are<sup>4,5,7</sup>

$$\dot{\theta} = JS (\cos\theta \nabla^2 \phi - 2 \sin\theta \nabla\theta \cdot \nabla\phi) \quad (4a)$$

$$\dot{\phi} \cos\theta = -\frac{1}{2} JS \{ [\delta (|\nabla\theta|^2 - 4) + |\nabla\phi|^2] \sin 2\theta + 2(1 - \delta \cos^2\theta) \nabla^2 \theta \} \quad (4b)$$

where  $\delta = 1 - \lambda$ . Using polar coordinates  $(r, \varphi)$ , and assuming a

